


LISTING OF CLAIMS

We claim:

1-30 (canceled)

31. (new) A compound selected from the group of compounds represented by Formula (I)



wherein:

X is H₂ or CH₂;

R¹ and R² are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl;

R³ and R⁴ are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl; or R³ and R⁴ together with C25 form a (C₃-C₉)cycloalkyl or (C₃-C₉)cyclofluoroalkyl;

32. (new) The compound of Claim 31, wherein:

R³ and R⁴ are, independently of each other, a (C₁-C₄)alkyl or a (C₁-C₄)fluoroalkyl;

X is =CH₂.

33. (new) The compound of Claim 32, wherein:

R¹ and R² are methyl; and

R³ and R⁴ are, independently of each other, methyl, ethyl, trifluoromethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl.

34. (new) The compound of Claim 33, wherein R³ and R⁴ are methyl.

35. (new) The compound of Claim 33, wherein R³ and R⁴ are trifluoromethyl.

36. (new) The compound of Claim 31, wherein:

R^3 and R^4 are, independently of each other, a (C₁-C₄)alkyl or a

(C₁-C₄)fluoroalkyl;

X is H₂.

37. (new) The compound of Claim 36, wherein:


R^1 and R^2 are methyl; and

R^3 and R^4 are, independently of each other, methyl, ethyl, trifluoromethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl.

38. (new) The compound of Claim 37, wherein R^3 and R^4 are methyl.

39. (new) The compound of Claim 37, wherein R^3 and R^4 are trifluoromethyl.

40. (new) A method of treating osteoporosis via administration of a therapeutically effective amount of a compound of Formula (I)ⁱ




wherein:

X is hydrogen or =CH₂;

R^1 and R^2 are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl,;

R^3 and R^4 are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl, or R^3 and R^4 together with C25 form a (C₃-C₉)cycloalkyl or (C₃-C₉)cyclofluoroalkyl.

41. (new) A method of treating cancer via administration of a therapeutically effective amount of a compound of Formula (I)




wherein:

X is H₂ or CH₂;

R¹ and R² are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl;

R³ and R⁴ are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl, or R³ and R⁴ together with C25 form a (C₃-C₉)cycloalkyl or (C₃-C₉)cyclofluoroalkyl;

42. (new) A method of treating secondary hyperparathyroidism via administration of a therapeutically effective amount of a compound of Formula (I)



wherein:

X is H₂ or CH₂;

R¹ and R² are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl;

R³ and R⁴ are, independently of each other, a (C₁-C₄)alkyl or (C₁-C₄)fluoroalkyl, or R³ and R⁴ together with C25 form a (C₃-C₉)cycloalkyl or (C₃-C₉)cyclofluoroalkyl;

43. (new) A pharmaceutical composition comprising an effective amount of a compound of Claim 31.

* * * * *